



Seasonal affective disorder

Seasonal affective disorder is a mental health condition that is triggered by the changing of the seasons. This condition is a subtype of major depressive disorder and bipolar disorder. Major depressive disorder is characterized by prolonged sadness and a general lack of interest, while bipolar disorder is characterized by similar depressive episodes alternating with periods of abnormally high energy and activity (hypomania or mania). People with seasonal affective disorder have signs and symptoms of either major depressive disorder or bipolar disorder only during certain months of the year. Major depressive disorder is more common than bipolar disorder among people with seasonal affective disorder. This condition usually begins in a person's twenties or thirties.

The signs and symptoms that occur during depressive episodes in people with seasonal affective disorder are similar to those of major depressive disorder, including a loss of interest or enjoyment in activities, a decrease in energy, a depressed mood, and low self-esteem. In most people with seasonal affective disorder, depression and other features appear in the fall and winter months and subside in the spring and summer months. In these individuals, additional symptoms often include weight gain due to increased intake of carbohydrates and an increase in sleep (hypersomnia). Affected individuals with underlying bipolar disorder typically have alternating episodes of depression in the fall and winter months and mania during the spring and summer months.

In about 10 percent of people with seasonal affective disorder, the condition has the opposite seasonal pattern, occurring in the spring and summer months and stopping during the fall and winter months. These individuals usually have a loss of appetite and sleep, unlike those with symptoms in the fall and winter.

For those affected, it is estimated that symptoms of seasonal affective disorder are present during 40 percent of the year. In some individuals, seasonal affective disorder does not recur every year. Thirty to 50 percent of affected individuals do not show symptoms of the disorder in consecutive winters. In about 40 percent of individuals with seasonal affective disorder, depressive episodes continue after winter and do not alleviate in the summer months, leading to a change in diagnosis to either major depressive disorder or bipolar disorder.

Individuals with seasonal affective disorder tend to have another psychological condition, such as attention deficit hyperactivity disorder (ADHD), an eating disorder, anxiety disorder, or panic disorder.

Frequency

Seasonal affective disorder occurs in 0.5 to 3 percent of individuals in the general population; it affects 10 to 20 percent of people with major depressive disorder and about 25 percent of people with bipolar disorder.

Some individuals have a condition known as subsyndromal seasonal affective disorder or seasonality, which is more common than seasonal affective disorder. These individuals have only mild changes in mood that correspond with the changes in seasons.

Genetic Changes

The causes of seasonal affective disorder are complex. A shortage of sunlight contributes to the development of the disorder in the fall and winter months, and too much sunlight is associated with seasonal affective disorder in the spring and summer months. Affected individuals seem to have disrupted daily (circadian) rhythms, such as the sleep-wake cycle, which are normally regulated to match the night-day cycle. These individuals cannot alter their sleep-wake cycle to match the night-day cycle of the winter months, resulting in changes in sleep, mood, and behavior. Those with seasonal affective disorder in summer have difficulty adjusting to the increased daylight hours. It is likely that affected individuals are genetically predisposed to being unable to adjust their circadian rhythms. Studies have identified variants in multiple genes that are associated with developing seasonal affective disorder. However, few of these variants have been verified in multiple studies.

Most genes that have been implicated in seasonal affective disorder are active (expressed) in the brain, where they are involved in the regulation of circadian rhythms. Some of these genes play a role in the expression of certain genes at specific times during the day or night, which helps set circadian rhythms. Others are involved in the production of chemical messengers in the brain known as neurotransmitters, specifically dopamine and serotonin. Dopamine can be converted into hormones that control blood pressure and body temperature, which change depending on the time of day. Serotonin can be converted into a hormone called melatonin, which signals to the brain that it is time to sleep.

Abnormal regulation of circadian rhythms likely contribute to an individual's inability to match their circadian rhythms to changes in seasons, resulting in changes in sleep, mood, and behavior as daylight hours change. However, little is known about what causes certain individuals to be more sensitive to the changing of the seasons than others.

Inheritance Pattern

The inheritance pattern of seasonal affective disorder is unclear. Overall, the risk of developing this condition is greater for first-degree relatives (such as parents or siblings) of affected individuals compared to the general public. Studies have found that

approximately 15 percent of individuals with seasonal affective disorders have a first-degree relative who also has the condition.

Many individuals with seasonal affective disorder have relatives with other mood or psychological disorders (such as major depressive disorder or schizophrenia). Studies show that 25 to 67 percent of people with seasonal affective disorder have one or more relatives with such a disorder. These other disorders may run in families in part because they share some genetic risk factors with seasonal affective disorder. However, these conditions are relatively common in the general population, and so they may occur in multiple people in a family by chance.

Other Names for This Condition

- affective disorder, seasonal
- depression in a seasonal pattern
- depression; seasonal
- major depressive disorder with a seasonal pattern
- SAD
- seasonal depression
- seasonal mood disorder

Diagnosis & Management

Formal Diagnostic Criteria

- U.S Preventive Services Task Force: Final Recommendation Statement: Depression Screening in Adults
<https://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/depression-in-adults-screening1>

Formal Treatment/Management Guidelines

- Agency for Healthcare Research and Quality: Practice Guideline for the Treatment of Patients with Major Depressive Disorder
<https://www.guideline.gov/summaries/summary/24158/practice-guideline-for-the-treatment-of-patients-with-major-depressive-disorder-third-edition>

Other Diagnosis and Management Resources

- American Psychological Association
<http://www.apa.org/helpcenter/seasonal-affective-disorder.aspx>
- American Psychological Association: Seasonal Affective Disorder Sufferers Have More Than Just Winter Blues
<http://www.apa.org/news/press/releases/2013/02/seasonal-disorder.aspx>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Health Topic: Seasonal Affective Disorder
<https://medlineplus.gov/seasonalaffectivedisorder.html>

Additional NIH Resources

- National Institute of Mental Health
<https://www.nimh.nih.gov/health/topics/seasonal-affective-disorder/index.shtml>
- NIH News in Health: Beat the Winter Blues
<https://newsinhealth.nih.gov/2013/01/beat-winter-blues>

Educational Resources

- Cleveland Clinic
<https://my.clevelandclinic.org/health/diseases/9293-seasonal-depression>
- Johns Hopkins Medicine
https://www.hopkinsmedicine.org/healthlibrary/conditions/adult/mental_health_disorders/seasonal_affective_disorder_85,p00755
- Merck Manual Consumer Version: Bipolar Disorder
<https://www.merckmanuals.com/home/mental-health-disorders/mood-disorders/bipolar-disorder>
- Merck Manual Consumer Version: Depression
<https://www.merckmanuals.com/home/mental-health-disorders/mood-disorders/depression>
- National Alliance on Mental Illness: Major Depressive Disorder with a Seasonal Pattern
https://www.nami.org/NAMI/media/NAMI-Media/Images/FactSheets/Major-Depressive-Disorder-with-a-Seasonal-Pattern_factsheet.pdf

- TeensHealth from Nemours
<http://kidshealth.org/en/teens/sad.html>
- University of Michigan
<https://www.uofmhealth.org/health-library/hw169553>
- University of Rochester Medical Center
<https://www.urmc.rochester.edu/encyclopedia/content.aspx?contenttypeid=85&contentid=P00755>

Patient Support and Advocacy Resources

- Anxiety and Depression Association of America
<https://adaa.org/>
- Depression and Bipolar Support Alliance
<http://www.dbsalliance.org/site/PageServer?pagename=home>
- International Bipolar Foundation
<http://ibpf.org/>
- National Alliance on Mental Illness
<https://www.nami.org/>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22seasonal+affective+disorder%22>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Seasonal+Affective+Disorder%5BMAJR%5D%29+AND+%28seasonal+affective+disorder%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- MAJOR DEPRESSIVE DISORDER
<http://omim.org/entry/608516>

Sources for This Summary

- Albrecht U. Circadian clocks and mood-related behaviors. *Handb Exp Pharmacol*. 2013;(217): 227-39. doi: 10.1007/978-3-642-25950-0_9. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23604481>
- Danilenko KV, Levitan RD. Seasonal affective disorder. *Handb Clin Neurol*. 2012;106:279-89. doi: 10.1016/B978-0-444-52002-9.00017-6. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22608628>

- Etain B, Milhiet V, Bellivier F, Leboyer M. Genetics of circadian rhythms and mood spectrum disorders. *Eur Neuropsychopharmacol*. 2011 Sep;21 Suppl 4:S676-82. doi: 10.1016/j.euroneuro.2011.07.007. Epub 2011 Aug 10. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21835597>
- Geoffroy PA, Bellivier F, Scott J, Etain B. Seasonality and bipolar disorder: a systematic review, from admission rates to seasonality of symptoms. *J Affect Disord*. 2014 Oct;168:210-23. doi: 10.1016/j.jad.2014.07.002. Epub 2014 Jul 10. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25063960>
- McClung CA. Circadian rhythms and mood regulation: insights from pre-clinical models. *Eur Neuropsychopharmacol*. 2011 Sep;21 Suppl 4:S683-93. doi: 10.1016/j.euroneuro.2011.07.008. Epub 2011 Aug 11. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21835596>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3179573/>
- Partonen T. Clock gene variants in mood and anxiety disorders. *J Neural Transm (Vienna)*. 2012 Oct;119(10):1133-45. Epub 2012 Apr 27. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22538398>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/seasonal-affective-disorder>

Reviewed: December 2017

Published: March 20, 2018

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services